

The quantum number of *Color*, colored quarks and QCD

V. Matveev , A. Tavkhelidze

(INR, RAS)

Thirteenth Lomonosov Conference on Elementary Particle Physics,
MSU, August **23 – 29**, 2007

Content

- **Introduction**
- **The quantum number of color and colored quarks**
- **Dynamic quark models of hadrons composed of quasifree colored quarks**
 - a) **Hadron form factors**
 - b) **The model of quasifree quarks and the laws of scaling at high energies**
 - c) **Quark counting formulae**

Summary

Introduction

At present, the dominant point of view is that all physical phenomena and processes, both terrestrial and cosmological, are governed by three fundamental forces: gravitational, electroweak and chromodynamic. The color charge serves as the source of chromodynamic forces.

In this talk we shall expose the main stages of the early development of the hypothesis of the quantum number of color and of colored quarks, put forward under the ideological influence of and in collaboration with N. Bogolubov at the JINR Laboratory of Theoretical Physics. In these works, the concept of color, colored quarks, was introduced for the first time, and a dynamical description of hadrons was given within the framework of the model of quasifree colored quarks.

Introduction of the quantum number of color permitted to treat colored quarks as real physical objects, constituents of matter. Further, from the color $SU(3)$ symmetry, the Yang-Mills principle of local invariance and quantization of chromodynamic fields gave rise to quantum chromodynamics – the modern theory of strong interactions.

The quantum number of color and colored quarks

In 1964, when the hypothesis of quarks was put forward by Gell-Mann and Zweig, quarks were only considered to be mathematical objects, in terms of which it was possible, in a most simple and elegant way, to describe the properties, already revealed by that time, of the approximate unitary SU(3)-symmetry of strong interactions. At the beginning, these particles, exhibiting fractional charges and not observable in a free state, were not attributed the necessary physical interpretation.

First of all, making up hadrons of quarks, possessing spin $\frac{1}{2}$, led to a contradiction with the Pauli principle and the Fermi-Dirac statistics for systems composed of particles of semiinteger spin.

The problem of the quark statistics was not, however, the sole obstacle in the path of theory. No answer existed to the following question: why were only systems consisting of three quarks and quark-antiquark pairs realized in Nature, and why were there no indications of the existence of other multiquark states?

Especially important was the issue of the possible existence of quarks in a free state (the problem of quark confinement).

In 1965, analysis of these problems led N.Bogolubov, B.Struminsky and A.Tavkhelidze, as well as I.Nambu and M.Hana, and I.Miyamoto to the cardinal idea of quarks exhibiting a new, hitherto unknown, quantum number subsequently termed **color**.

From the very beginning creation of the relativistically invariant dynamical quark model of hadrons was based, first of all, on the assumption of quarks representing real physical objects determining the structure of hadrons.

To make it possible for quarks to be considered fundamental physical particles, the hypothesis was Proposed by three authors (Bogolubov, Struminsky, Tavkhelidze – 1965, January) that the quarks, should possess an **additional quantum number**, and that quarks of each kind may exist in three (unitary) equivalent states

$$\mathbf{q} \equiv (q^1, q^2, q^3)$$

differing in values of the new quantum number, subsequently termed **color**. Since at the time, when the new quantum number was introduced, only three kinds of quarks were known – (u,d,s), the quark model with an additional quantum number was termed the three-triplet model. Since the new quantum number is termed color, colored quarks may be in three equivalent states, such as, for example, **red, blue and green**.

With introduction of the new quantum number, color, the question naturally arised of the possible appearance of hadrons possessing color, which, however, have not been observed. From the assumption that colored quarks are physical objects, while the hadron world is degenerate in the new quantum number, or it is colorless, **it followed that solutions of the dynamic equations for baryons and mesons in the s-state should be neutral in the color quantum numbers.**

From the requirement that baryons be colorless, the wave function of the observed baryon family in the ground state, described by the totally symmetric 56-component tensor $\Phi_{abc}(x_1, x_2, x_3)$ in the approximation of spin-unitary symmetry, was assumed to be totally antisymmetric in the color variables of the three constituent quarks,

$$\Psi_{ABC}(x_1, x_2, x_3) = 1/\sqrt{6} \varepsilon_{\alpha\beta\gamma} \Phi_{abc}(x_1, x_2, x_3), \quad (\alpha, \beta, \gamma = 1, 2, 3),$$

where

$$A=(a, \alpha), \quad B=(b, \beta), \quad C=(c, \gamma),$$

a,b,c are unitary quantum numbers, α, β, γ are color quantum numbers, $\varepsilon_{\alpha\beta\gamma}$ is totally antisymmetric tensor.

Hence it is evident, that the Pauli principle holds valid for colored quarks, and that they satisfy the Fermi-Dirac statistics, so they can be considered real fundamental constituents of matter.

From the requirement that mesons, composed of a colored quark and an antiquark, be neutral, or colorless, the meson wave function is chosen in the form

$$\Psi_A^B(x_1, x_2) = 1/\sqrt{3} \delta_{\alpha}^{\beta} \Phi_a^b(x_1, x_2),$$

where a and b are unitary indices, δ_{α}^{β} is Kronecker symbol.

The choice of baryon and meson wave functions, proposed above, leads to the conclusion that the known mesons and baryons are composed of colored quarks and antiquarks as follows:

$$\bar{q}^\alpha(1)q_\alpha(2) \quad \text{– mesons;}$$

$$\varepsilon^{\alpha\beta\gamma}q_\alpha(1)q_\beta(2)q_\gamma(3) \quad \text{– baryons.}$$

Subsequently, the requirement that the world of hadrons are neutral led to the discovery of the SU(3) color symmetry group.

It is to be noted, here, that in his talk delivered to the conference “Symmetry principles at high energies”, held in Coral Gables (1965), Nambu was the first, on the basis of SU(3) symmetry with respect to the new quantum number (color), to deal with eight vector fields, carriers of the interaction between quarks, which were the prototype of the quantum-chromodynamic gluon fields.

Dynamic quark models of hadrons composed of quasifree colored quarks

a) Hadron form factors

The introduction of colored quarks, representing physical fundamental particles, paved the way for the dynamic description of hadrons.

The main obstacle, here, was the absence of quarks in a free state. Although it was evident that the issue of confinement could be ultimately settled only by experiments, a series of attempts was undertaken to provide a logically non-contradictory explanation for the “eternal confinement” of quarks inside hadrons. Thus, for example, P. Bogolubov (1968) proposed the “quark-bag” model known as the **Dubna bag**. Later (1974), the idea of a quark bag underwent development at MIT, and the resulting model is known as the **MIT bag**.

The dynamic relativistic quasi-free quark model, the development of which was initiated in Dubna (1965 – January, April), was based on the assumption of quarks being extremely heavy objects bound in hadrons by enormous scalar forces, that on the one hand provide for the large quark mass defect in hadrons and on the other hand impedes their leaving the hadron.

In this model baryons Φ_{ABC} and mesons Φ_A^B are represented by a superposition of all admissible states over the quantum numbers (A,B,C), satisfying the requirements of SU(6) symmetry, of quark statistics and of hadron neutrality in the color quantum number.

The dynamic composite quasifree quark model has made possible the systematic description of both the statically observed characteristics of hadrons (μ , g_A/g_V etc.) and their form factors. Weak and electromagnetic interactions were introduced in a minimal manner,

$$i\partial_\mu \rightarrow i\partial_\mu + \begin{cases} eA_\mu & \text{- electromagnetic interaction} \\ G \tau^\pm \gamma_5 l_\mu^\pm & \text{- weak interaction} \end{cases}$$

where A_μ is the electromagnetic potential, l_μ^\pm represent charged lepton weak currents, G is the Fermi weak interaction constant.

For the ratio g_A/g_V (of the axial and vector weak interaction constants) and for the magnetic moment of the proton we obtain

$$g_A/g_V \cong -5/3 (1-2\delta),$$

$$\mu_p \cong 3/(2M_p)(1-\delta),$$

$$\delta = \langle p | L_z | p \rangle$$


where L_z is the orbital momentum of a quark bound in the nucleon with the projection of its total angular momentum equal to $1/2$; δ characterizes the magnitude of relativistic corrections and amounts to $\delta \sim 1/6$, the resulting correction for the ratio g_A/g_V is of the order of 30%. This example shows how significant could be the relativistic corrections..

The dynamic quark model has permitted to explain the lepton decays of pseudoscalar π^- and K-mesons and, also, the electromagnetic decays of the vector mesons into electron-positron pairs as annihilation of quark-antiquark pairs bound in the mesons.

Analysis of the data on the widths of these decays resulted in a conclusion on the dependence of the scales of distances (effective sizes) on the quantum numbers of a bound system. For example

$$\frac{|\Psi_k(0)|^2}{|\Psi_\pi(0)|^2} \cong \frac{m_k}{m_\pi}.$$

In the case of the decay $\pi^0 \rightarrow 2 \gamma$,



determined by the triangular anomaly of the axial current (J.Bell, R.Jackiw 1969; S.Adler 1970), the **annihilation model points to the width of this decay being proportional to the number of different quark colors.**
(Matveev, Struminsky, Tavkhelidze - 1965).

b) The model of quasifree quarks and the laws of scaling at high energies

Experiments, in which inclusive reactions were studied at high energies and momentum transfers, and the scaling regularities revealed, have given an impetus to further development of the theory of hadron quark structure.


(Inclusive reactions were first introduced and studied theoretically by Logunov, Nguen Van Hieu and Mestvirishvili (1967); reported by Logunov at the Rochester conference (1967)).

Here, of essential significance was the investigation of deep inelastic processes in the inclusive scattering of electrons of nucleons, performed at the Stanford center, which in 1968 resulted in observation of the scaling properties - Bjorken scaling - indicating the existence of a "rigid" pointlike nucleonic structure.

In 1969, on the basis of the quasi-free quark model, the assumption was put forward by Matveev, Muradyan and Tavkhelidze that the scaling properties of electron-nucleon interaction processes, revealed in experiments, are common for all deep-inelastic lepton-hadron processes and that they can be derived in a model-independent manner on the basis of the automodelling principle, or the principle of self-similarity.

The essence of the self-similarity principle consists in the assumption that in the asymptotic limit of high energies and large momentum transfers form factors and other measurable quantities of deep-inelastic processes are independent of any dimensional parameters (such as particle masses, the strong interaction radius etc.), which may set the scale of measurement of lengths or momenta. **Thus, the form factors of deep-inelastic processes turn out to be homogeneous functions of relativistically invariant kinematic variables, the degree of homogeneity of which is determined by analysis of the dimensionality (it is a key property of conformal invariant theories).**

Application of the self-similarity principle for establishing the asymptotic behaviour of the form factors $W_1(q^2, \nu)$ and $W_2(q^2, \nu)$ of deep-inelastic scattering of electrons on protons in the Bjorken region




$$|q^2| \rightarrow \infty, \quad \nu = 2P \cdot q \rightarrow \infty,$$

$$\xi = -\frac{q^2}{2\nu} = \text{const}$$

results in the Bjorken asymptotic formulae derived in 1968 on the basis of certain assumptions

$$W_1(q^2, \nu) \rightarrow f_1(\xi), \quad \nu W_2(q^2, \nu) \rightarrow f_2(\xi) .$$

Application of the self-similarity principle resulted in the scaling law, being found for the first time, that describes the mass spectrum of muon pairs, produced in inclusive proton collisions,



$$\frac{d\sigma}{dM^2} \sim \frac{1}{M^4} \psi\left(\frac{M}{E}\right)$$

(Matveev, Muradyan, Tavkhelidze – 1969)

where M is the effective mass of the muon pair, and E is the initial energy of the colliding particles. (Later, this process was called Drell—Yan process).

Experimental studies of this process, initiated in 1970 by the group of L.Lederman at Brookhaven, confirmed this scaling law, and it was precisely in these processes that a new class of hadrons, the J/ψ particles, was subsequently observed.

c) Quark counting formulae

In the case of binary hadron reactions $a + b \rightarrow c + d$ at high energies s and momentum transfers t , application of the self-similarity principle yields for the differential cross section the following formula of the quark counting

$$\left(\frac{d\sigma}{dt} \right)_{a+b \rightarrow c+d} \sim \left(\frac{1}{s} \right)^{n-2} f\left(\frac{t}{s} \right),$$

$$n = n_a + n_b + n_c + n_d$$

(Matveev, Muradyan, Tavkhelidze – 1973)

where n is the total number of quarks belonging to the particles participating in the reactions. In the case, when particle b , for instance, is a lepton, then $n_b = 1$, and one obtains the asymptotic formula for the baryon form factor.

The function $f(t/s)$ depends only on the relation between large kinematic variables and is itself a dimensional quantity. Thus, the asymptotic power law points to factorization of the effects of large and small distances.

Summary

We described the scaling properties of elementary particle interaction processes, observed experimentally at high energies and large momentum transfers, on the basis of the self-similarity principle.

At the same time the question arises concerning the extent to which scaling invariant behaviour is consistent with the main requirements of local quantum field theory.

In the case of deep-inelastic electron scattering on nucleons, these problems were investigated by Bogolubov, Vladimirov and Tavkhelidze (1972) for form factors, which in the Bjorken region have the asymptote

$$W_1(q^2, \nu) \rightarrow f_1(\nu/q^2), \quad \nu W_2(q^2, \nu) \rightarrow f_2(\nu/q^2).$$

For the weight functions of these form factors in the Jost-Lehman-Dyson representation sufficient conditions were found that guarantee Bjorken scaling. In the case of free field the weight functions automatically satisfy these restrictions, which is precisely what provides for Bjorken scaling in the quasi-free quark model.

Note, that the subsequent discovery (1973) by Gross, Wilczek and Politzer of the phenomenon of asymptotic freedom in QCD of an invariant charge, introduced by Bogolubov and Shirkov in the renormgroup theory, is an essentially important step for substantiation of the picture of quasi-free quarks in hadrons.

.

In a number of important works of the last years (t'Hooft, Maldacena, Polchinski, Polyakov, Witten) there was suggested an impressive non-perturbative derivation of the asymptotic power laws (the quark counting for form factors and the exclusive scattering cross sections of hadrons) in the framework of the conformal versions of QCD dual to the string theory.

In this talk we have not touch upon the possibility of parafermionic character of quarks, suggested by Greenberg in 1964, as we assume that this problem shall be mentioned here by Prof. O.W.Greenberg himself.

It can be noted that many of the priority works, presented in this talk, were only published as preprints or, sometimes, in the proceedings of international conferences, in accordance with the opportunities that existed at that time.

The references to the original paper mentioned in the talk are presented “PARTICLES AND NUCLEUS” 2006 v. 37 P.3 576 – 595 V.Matveev, A.Tavkhelidze THE QUANTUM NUMBER COLOR QUARKS AND QCD (Dedicated to the 40th anniversary of the discovery of color.